

Psychological Science

<http://pss.sagepub.com/>

Meditation Increases Compassionate Responses to Suffering

Paul Condon, Gaëlle Desbordes, Willa B. Miller and David DeSteno

Psychological Science published online 21 August 2013

DOI: 10.1177/0956797613485603

The online version of this article can be found at:

<http://pss.sagepub.com/content/early/2013/08/21/0956797613485603>

Published by:



<http://www.sagepublications.com>

On behalf of:



[Association for Psychological Science](http://www.sagepublications.com)

Additional services and information for *Psychological Science* can be found at:

Email Alerts: <http://pss.sagepub.com/cgi/alerts>

Subscriptions: <http://pss.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [OnlineFirst Version of Record](#) - Aug 21, 2013

[What is This?](#)

Meditation Increases Compassionate Responses to Suffering

Paul Condon¹, Gaëlle Desbordes², Willa B. Miller³,
and David DeSteno¹

¹Department of Psychology, Northeastern University; ²Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Harvard Medical School; and ³Department of Religion, Harvard University

Received 1/3/13; Revision accepted 3/17/13

Contemplative science has documented a plethora of intrapersonal benefits stemming from meditation, including increases in gray matter density (Hölzel, Carmody, et al., 2011), positive affect (Moyer et al., 2011), and improvement in various mental-health outcomes (Hölzel, Lazar, et al., 2011). Strikingly, however, much less is known about the interpersonal impact of meditation. Although Buddhist teachings suggest that increases in compassionate responding should be a primary outcome of meditation (Davidson & Harrington, 2002), little scientific evidence supports this conjecture. Even as scientists have begun to examine the effects of meditation on prosocial action, the conclusions that can be drawn with respect to compassion have been limited by designs that lack real-time person-to-person interactions centered on suffering. Previous work, for example, has utilized meditators' self-reported intentions and motivations to behave in supportive manners toward other individuals (e.g., Fredrickson, Cohn, Coffey, Pek, & Finkel, 2008) and computer-based economic games requiring cooperation (e.g., Leiberg, Klimecki, & Singer, 2011; Weng et al., 2013) to assess altruistic action. Such methods have suggested that meditation may increase generalized prosocial responding, but have not clearly and objectively gauged responses meant solely to mitigate the suffering of other individuals.

To address this gap, we utilized a design in which individuals were confronted with a person in pain in an ecologically valid setting. If, as suggested by Buddhist theorizing, meditation enhances compassionate responding, participants who have completed a brief meditation course should act to relieve such a person's suffering more frequently than those who have not completed the course.

Method

The final set of participants comprised 39 individuals (29 female, 10 male; mean age = 25.23 years, $SD = 4.66$)

recruited from the Greater Boston community for an 8-week study on meditation. (See the Supplemental Material available online for recruitment procedures.) Individuals were randomly assigned either to complete meditation classes or to be in a waiting-list control group. Those assigned to the meditation condition were further randomly subdivided to receive one of two protocols: mindfulness or compassion meditation. We utilized two separate meditation protocols both to enhance generalizability and to ensure that any resulting effects of meditation on behavior could not be attributed to demand characteristics. Although techniques to focus and calm the mind were taught in both protocols, direct discussion of compassion and the suffering of other people occurred only in compassion meditation. (See the Supplemental Material for meditation protocols.)

Meditation classes were held in a nondenominational venue dedicated to spiritual activities (e.g., prayer, meditation, yoga). A Tibetan Buddhist lama (author W. M.) with 20-plus years of meditation experience conducted both courses. The classes were taught in a secular format featuring 60 min of instruction, 30 min of practice, and 30 min for discussion; classes were held once a week for 8 weeks. Participants also received 20-min audio-guided meditations to complete independently outside of class. Participants reported their daily use of the audio recordings each week. Participants received \$60 for their participation.¹

Following 8 weeks of meditation practice or approximately 8 weeks after initial recruitment to the waiting list,

Corresponding Authors:

Paul Condon, Department of Psychology, 360 Huntington Ave.,
Northeastern University, Boston, MA 02115
E-mail: p.condon@neu.edu

David DeSteno, Department of Psychology, 360 Huntington Ave.,
Northeastern University, Boston, MA 02115
E-mail: d.desteno@gmail.com

Psychological Science
XX(X) 1–3
© The Author(s) 2013
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0956797613485603
pss.sagepub.com


participants were scheduled to come to the lab under the guise of completing tests of cognitive ability. To obtain a naturalistic measure of responses to suffering, we utilized confederates to construct a test situation outside the laboratory. All confederates were blind both to the hypothesis being tested and to each participant's experimental condition. Prior to a participant's arrival, two female confederates sat in a designated waiting area possessing three chairs. Upon arriving at the waiting area, the participant sat in the remaining chair. After the participant had been sitting for 1 min, a third female confederate, who played the role of the "sufferer," appeared around the corner with crutches and a walking boot. The sufferer, who visibly winced while walking, stopped just as she arrived at the chairs. She then looked at her cell phone, audibly sighed in discomfort, and leaned back against a wall.

To assess compassionate responding, we measured whether the true participant offered his or her seat to the sufferer to relieve her pain. Via text message, one of the sitting confederates surreptitiously notified the experimenter, who was waiting out of sight, whether and when the participant offered the seat to the sufferer. If 2 min passed and the participant had not given up his or her seat, the trial was ended and coded as a nonhelping response. The experimenter then entered the waiting area, greeted the participant, and escorted him or her to the lab to complete a series of measures unrelated to the goals of the present analysis.

Results and Discussion

As predicted, meditation directly enhanced compassionate responding. Meditators offered their seats to the sufferer more frequently than did nonmeditators from the waiting-list control group, $\chi^2(1, N = 39) = 5.13, p = .02, \phi = .36$ (see Table 1). This enhanced prosocial responding did not differ as a function of meditation protocol; participants practicing mindfulness meditation were as likely to aid the sufferer as were those practicing compassion meditation (see the Supplemental Material for analysis).² That 8 weeks of meditation resulted in such a large effect—increasing the odds of acting to relieve another person's pain by more than 5 times (odds

Table 1. Observed and Expected Frequencies of Helping Behavior Across Conditions

Outcome	Meditation training		Waiting-list control	
	Observed	Expected	Observed	Expected
No help	10	13.3	16	12.7
Help	10	6.7	3	6.3

ratio = 5.33)—is all the more striking given that the helping occurred in a social context whose features should attenuate such behavior. The simple presence of the two confederates and their total disregard for the pain of the sufferer constitutes a classic bystander manipulation in which both diffusion of responsibility and norms suggesting an acceptance of nonintervention are heightened (cf. Darley & Latané, 1968).

Additional work will be needed to isolate the specific causal mechanism for the observed effect more narrowly, as several meditation-induced mediators (e.g., heightened awareness, increased perspective taking) stand as possible candidates (cf. Hölzel, Lazar, et al., 2011). Nonetheless, this study is the first to clearly show the power of meditation to increase compassionate responding to suffering, even in the face of social pressures to avoid so doing. In turn, it provides scientific credence to ancient Buddhist teachings that meditation increases spontaneous compassionate behavior.

Author Contributions

P. Condon, D. DeSteno, G. Desbordes, and W. B. Miller designed the experiments. P. Condon and W. B. Miller conducted the research. P. Condon and D. DeSteno conducted the analyses and wrote the manuscript.

Declaration of Conflicting Interests

The authors declared that they had no conflicts of interest with respect to their authorship or the publication of this article.

Funding

We acknowledge support from a Mind and Life Institute Francisco J. Varela award. Any views, findings, conclusions, or recommendations expressed in this publication do not necessarily reflect those of the Mind and Life Institute.

Supplemental Material

Additional supporting information may be found at <http://pss.sagepub.com/content/by/supplemental-data>

Notes

1. Participants in the meditation condition earned an additional \$20 and entry into a \$100 raffle for completing weekly logs.
2. Additional analyses revealed that gender did not affect rates of helping behavior. Also of import, analyses of self-reported social networks demonstrated that social-network size neither increased as a result of participation in the meditation classes nor significantly differed between the meditation and control groups, suggesting that increases in social capital from participating in a group activity could not account for the central finding that meditation increased compassionate responding (see the Supplemental Material for these and additional results).

References

- Darley, J. M., & Latané, B. (1968). Bystander intervention in emergencies: Diffusion of responsibility. *Journal of Personality and Social Psychology*, *8*, 377–383.
- Davidson, R. J., & Harrington, A. (2002). Visions of compassion: Western scientists and Tibetan Buddhists examine human nature. New York, NY: Oxford University Press.
- Fredrickson, B. L., Cohn, M. A., Coffey, K. A., Pek, J., & Finkel, S. M. (2008). Open hearts build lives: Positive emotions, induced through loving-kindness meditation, build consequential personal resources. *Journal of Personality and Social Psychology*, *95*, 1045–1062.
- Hölzel, B. K., Carmody, J., Vangel, M., Congleton, C., Yerramsetti, S. M., Gard, T., & Lazar, S. W. (2011). Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Research: Neuroimaging*, *191*, 36–43.
- Hölzel, B. K., Lazar, S. W., Gard, T., Schuman-Olivier, Z., Vago, D. R., & Ott, U. (2011). How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspectives on Psychological Science*, *6*, 537–559.
- Leiberg, S., Klimecki, O., & Singer, T. (2011). Short-term compassion training increases prosocial behavior in a newly developed prosocial game. *PLoS ONE*, *6*, e17798. Retrieved from <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0017798>
- Moyer, C. A., Donnelly, M. P. W., Anderson, J. C., Valek, K. C., Huckaby, S. J., Widerholt, D. A., . . . Rice, B. L. (2011). Frontal electroencephalographic asymmetry associated with positive emotion is produced by very brief meditation training. *Psychological Science*, *22*, 1277–1279.
- Weng, H. Y., Fox, A. S., Shackman, A. J., Stodola, D. E., Caldwell, J. Z. K., Olson, M. C., . . . Davidson, R. J. (2013). Compassion training alters altruism and neural responses to suffering. *Psychological Science*, *24*, 1171–1180.

Supplementary Online Material (word count: 977)

Participants

During recruitment, all participants self-reported little to no experience with any type of meditation experience and none reported completing any previous meditation course or meditation retreat. All participants passed a telephone-administered version of the Mini-Mental State Examination as indicated by a score greater than or equal to 21 (Newkirk et al., 2004).

Sixty-seven individuals were initially recruited for the study. Twenty-six dropped out prior to completion of the training portion. Two others were removed by experimenters: one who did not take a chair in the waiting area upon arrival and one who expressed suspicion about the study. The final sample consisted of 39 individuals (20 meditators, and 19 waitlist controls).

Supplementary Analyses

Amount of Training

On average, meditating participants attended 6.60 ($SD=0.50$) instruction sessions and reported completing independent 20-minute intervals of meditation 3.74 ($SD=1.04$) times per week.

Helping Behavior by Meditation Class

Those participants who completed the mindfulness- and compassion-based meditation did not differ in frequency of helping behavior, $\chi^2(1)=0.20, p>.65$ (see Table S1).

Gender

Gender differences in helping behavior have been reported in numerous studies, although the direction of these differences vary depending on context. Males help more frequently in some contexts (e.g., short-term encounters), but females help more in others (e.g., long-term close relationships; for a review, see Eagly & Crowley, 1986). In the current study, we found that participant gender did not affect frequency of compassionate responding, $f_{\text{females}}=10$ out of 29, $f_{\text{males}}=3$ out of 10, $\chi^2(1, N=39)=0.07, p>.79$. Furthermore, the composition of male and female participants in the meditation and control groups did not differ, $\chi^2(1)=0.41$.

Design Limitations: Control Group and Social Network Analysis

The nature of our design required that one group (i.e., meditators) came together for repeated classes, thereby creating a context that afforded interaction with other individuals participating in the study. The waitlist group had no such possibility of interacting with others due to participation in a structured class. Thus, although the control group was matched with respect to interest in meditation and desire to enroll in an 8-week course, its members did not engage in repeated interactions with meditation class members or an instructor. One resulting concern is that repeated interactions with fellow participants in the course may have produced social consequences that could account for increased levels of helping behavior (e.g., increased social resources). To rule out this possibility, we obtained a measure of the number of people that participants interacted with on a regular basis before and after training. At pre- and post-testing (i.e., eight weeks apart), participants received an email with a link to an online version of the Social Network Index (Cohen, Doyle, Skoner, Rabin, & Gwaltney, 1997). This survey asked participants to list the initials of every individual that they “interacted with in person or over the phone at least once every two weeks.” The survey prompted participants to list people in the following categories: *romantic partner, parents, partner’s parents, children, relatives, friends, classmates, co-workers, acquaintances, members of a spiritual group, members of a volunteer group, and members of unspecified groups*. Participants could list up to seven individuals for each category, with the exception of *friends* and *members of unspecified group* (up to ten individuals). Our primary interest concerned the total number of people with whom participants reported interacting. If the meditation classes increased social capital, we would expect to find that those in the meditation group reported an increase in their number of relationships at post-testing, relative to the wait-list group.

A 2 (time: pre, post) X 2 (group: meditators, wait-list) repeated measures ANOVA, with time as the repeated factor, revealed no main effect of group on social capital ($M_{\text{meditators}}=11.58; SD_{\text{meditators}}=4.53; M_{\text{wait-list}}=12.35 SD_{\text{wait-list}}=5.07$), $F(1,31)=0.21, p>.65$, and no effect of time ($M_{\text{pre-test}}=12.18; SD_{\text{pre-test}}=5.60; M_{\text{post-test}}=11.58 SD_{\text{post-test}}=5.10$), $F(1,31)=0.83, p>.37$. There was no interaction, $F(1,31)=1.40, p>.24$ (note that only 13 participants from the WL control group provided responses to the SNI). In sum, participants

in the meditation group did not experience a growth in their social network as a function of participating in an organized class. Thus, the experience of participating in a group activity is unlikely to account for our central finding, at least as stemming from increases in social capital.

It could also be argued, however, that exposure to a caring or charismatic instructor might have exerted an influence, principally through modeling. That is, meditation in and of itself covaried with exposure to an instructor in our design. Although certainly possible, we believe that such a factor is unlikely to account for our findings in the present case. If it were the influence of exposure to a caring instructor, we would expect to find a greater effect of meditation on prosocial behavior among

participants following the compassion meditation protocol, where actual discussion of the virtue of relieving the suffering of others was discussed by the instructor. In the mindfulness meditation condition, no such discussions occurred; all instruction centered on techniques involved in centering attention (e.g., breathing), and consequently provided no opportunity for prosocial behavior to be modeled on or directly influenced by the instructor's explicit or implicit goals. Nonetheless, having established the basic phenomenon, we believe it fruitful for future investigation to aim toward disambiguating the exact mechanisms associated with contemplative training that may underlie its enhancement of prosocial behavior, as well as individual susceptibility to such enhancement.

References for Supplementary Material

- Cohen, S., Doyle, W.J., Skoner, D.P., Rabin, B.S., Gwaltney, J.M. (1997). Social ties and susceptibility to the common cold. *Journal of the American Medical Association*, 277, 1940-1944.
- Eagly, A.H., & Crowley, M. (1986). Gender and helping behavior: A meta-analytic review of the social psychological literature. *Psychological Bulletin*, 100, 283-308.
- Newkirk, L.A., Kim, J.M., Thompson, J.M., Tinklenberg, J.R., Yesavage, J.A., & Taylor, J.L. (2004). Validation of a 26-point telephone version of the Mini-Mental State Examination. *Journal of Geriatric Psychiatry and Neurology*, 17, 81-87.

Acknowledgement

We thank Kevin Bickart for assistance with the social network analysis.

Supplementary Table 1.*Observed and expected frequencies of helping behavior by meditation group.*

Outcome	Mindfulness-group		Compassion-group	
	Observed	Expected	Observed	Expected
No help	5	4.5	5	5.5
Help	4	4.5	6	5.5

Note. $\chi^2(1)=0.20, p>.65$

Supplementary Table 2A.*Mindfulness-based training protocol.*

Week	Training components
1	Open awareness meditation Introduction of basic techniques (e.g., body-scan) for relaxing the body and monitoring the mind's natural tendency to wander from the object of attention (i.e., the body).
2	Mindfulness of a physical object Introduction and elaboration of practices for learning to calm the conceptually discursive mind for the purpose of attenuating involuntary thoughts. Stability of attention (i.e., on a physical object) is practiced with the goal of sustaining attention in a purposeful, non-judgmental manner for an extended period.
3	Mindfulness of the breathing with relaxation (I) Continuing practice of techniques designed to instill a deepening sense of physical and mental relaxation, stillness, and vigilance. When successful, involuntary thoughts subside and vividness of attention gradually increases. This gives rise to an overall sense of greater presence, calm, and equilibrium.
4	Mindfulness of the breathing with relaxation (II) Continuing practice from Week 3. Additional instruction focused on common impediments to meditation practice, including sleepiness, agitation, and boredom.
5	Settling the mind in its natural state (i. e., mindfulness of mental events) (I) Introduction of practices for further refining the meditator's metacognitive abilities, with the goal of attenuating the immediate and habitual absorption in one's thoughts that characterize most mental functioning. When successful, insight into the nature of the mind and its activities is achieved.
6	Settling the mind in its natural state (II) Continued practice with the goal of developing increased relaxation, stillness of awareness in the midst of mental activities, and vividness, together with heightened metacognitive abilities to observe mental states and processes without identifying with them.
7	Awareness of awareness (I) In this final technique, relaxation, stillness, and vividness of attention continue to be enhanced, leading to a perception of the process of becoming aware, as opposed to only perceiving the contents of awareness.
8	Mindfulness of awareness and thoughts Participants use the techniques they've learned from all previous weeks and apply them to concentration on the quality of immediate mental experience. Instead of using breath or body as an object, now mind itself becomes the meditation object. Particular attention will be paid to working with the arising and disappearing of thoughts.

Supplementary Table 2B.*Compassion-based training protocol.*

Week	Training components
1	Developing attention and stability of mind Introduction of basic meditation techniques for focusing attention for increasingly longer periods of time. <i>These techniques are included in the practice of all subsequent compassion meditation components.</i>
2	Baring witness to ones current life-stressors and difficulties Introduction of practice for noticing and reflecting on current life challenges, such as a difficult situation, event, or person. Participants learn to accept themselves in these moments and relax resistance to discomfort to remain and experience these states without judgment or a need to remedy it. When successful, this practice leads to the dissolution of discomfort.
3	Commune with other's life-stressors and difficulties Introduction of techniques to develop awareness of the commonality of one's own suffering. Participants learn to recognize that all others feel discomfort just as they do. Participants imagine themselves in a community of others experiencing the same discomfort.
4	Extending compassion outward Building on previous practice of bearing witness and communing with suffering, participants visualize breathing own and others suffering into the heart (inhalation) and dissolving all difficulties outward into a spacious sky (exhalation).
5	Release of suffering Introduction of practice for letting go of one's own and others' suffering into a state of open awareness.
6	Exchange with others – close target The culmination of previous instruction are specifically applied to a close other (i.e., a person that one cares about). Participants visualize the disappearance of another's suffering as they breath it inward and send compassion outward.
7	Exchange with others – neutral, unknown target and a difficult, annoying target Extending practice of exchange with others to neutral and difficult targets.
8	Exchange with others – all sentient beings Extending practice of exchange with others to all sentient beings.